

Children's Hospital and Health System Children's Community Health Plan Policy and Procedure

This policy applies to the following entity(s):

- | | |
|--|--|
| <input type="checkbox"/> CHW – Milwaukee | <input type="checkbox"/> CHW - Fox Valley |
| <input type="checkbox"/> CHHS Foundation | <input type="checkbox"/> CHW - Surgicenter |
| <input type="checkbox"/> CHW – Community Services Division | <input checked="" type="checkbox"/> Children's Community Health Plan |
| <input type="checkbox"/> Children's Medical Group - Primary Care | <input type="checkbox"/> Children's Specialty Group |
| <input type="checkbox"/> Children's Medical Group - Urgent Care | <input type="checkbox"/> CHHS Corporate Departments |

Medical Utilization Management Policy

SUBJECT: CONTINUOUS PASSIVE MOTION DEVICES

INCLUDED PRODUCT(S):

Medicaid

BadgerCare Plus

Care4Kids Program

Commercial

Together with CCHP

Marketplace

Together with CCHP

PURPOSE OR DESCRIPTION:

The purpose of this policy is to define criteria for the medically necessary use of Continuous Passive Motion devices (CPM).

POLICY:

Appropriate use of CPMs is limited to:

1. Post operative use after total knee replacement surgery (knee arthroplasty) AND
 - a. Applied within 48 hours of surgery
 - b. Duration of use is limited to 21 days after surgery

For all other uses, CCHP considers the current role to be uncertain. Based on review of existing evidence, there are currently no other clinical indications for when this technology is proven beneficial.

Effective: 5/16
Revised: 2/20
Reviewed: 10/21

Q: \CCHP Leadership\Utilization Management Medical Policies\APPROVED MEDICAL UM POLICIES\Continuous Passive Motion Devices Medical UM Policy
Developed by: CCHP Medical Directors and Executive Director Health Plan Clinical Services

REFERENCES

1. MCG Guideline ACG: A-0335 (AC), Continuous Passive Motion (CPM); MCG Health, Ambulatory Care 23rd Edition, Copyright © 2019 MCG Health, LLC
2. Dundar U, Toktas H, Cakir T, Evcik D, Kavuncu V. Continuous passive motion provides good pain control in patients with adhesive capsulitis. *International Journal of Rehabilitation Research* 2009;32(3):193-8. DOI: 10.1097/MRR.0b013e3283103aac.
3. Lobb R, Tumilty S, Claydon LS. A review of systematic reviews on anterior cruciate ligament reconstruction rehabilitation. *Physical Therapy in Sport* 2012;13(4):270-8. DOI: 10.1016/j.ptsp.2012.05.001.
4. Fazalare JA, Griesser MJ, Siston RA, Flanigan DC. The use of continuous passive motion following knee cartilage defect surgery: a systematic review. *Orthopedics* 2010;33(12):878. DOI: 10.3928/01477447-20101021-16.
5. Garofalo R, Conti M, Notarnicola A, Maradei L, Giardella A, Castagna A. Effects of one-month continuous passive motion after arthroscopic rotator cuff repair: results at 1-year follow-up of a prospective randomized study. *Musculoskeletal Surgery* 2010;94 Suppl 1:S79-83. DOI: 10.1007/s12306-010-0058-7.
6. Harvey LA, Brosseau L, Herbert RD. Continuous passive motion following total knee arthroplasty in people with arthritis. *Cochrane Database of Systematic Reviews* 2014, (verified by Cochrane 2014 Mar), Issue 2. Art. No.: CD004260. DOI: 10.1002/14651858.CD004260.pub3.
7. He ML, Xiao ZM, Lei M, Li TS, Wu H, Liao J. Continuous passive motion for preventing venous thromboembolism after total knee arthroplasty. *Cochrane Database of Systematic Reviews* 2014, Issue 7. Art. No.: CD008207. DOI: 10.1002/14651858.CD008207.pub3

Effective: 5/16
Revised: 2/20
Reviewed: 10/21

Q: \CCHP Leadership\Utilization Management Medical Policies\APPROVED MEDICAL UM POLICIES\Continuous Passive Motion Devices Medical UM Policy
Developed by: CCHP Medical Directors and Executive Director Health Plan Clinical Services